

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number  
**WO 03/008387 A1**

(51) International Patent Classification<sup>7</sup>: C07D 239/69,  
403/12, A61K 31/635, A61P 9/10, 9/12

(74) Agent: TURUN PATENTTITOIMISTO OY; P.O. Box  
99, FIN-20521 Turku (FI).

(21) International Application Number: PCT/FI02/00643

(22) International Filing Date: 22 July 2002 (22.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
20011560 20 July 2001 (20.07.2001) FI  
60/306,449 20 July 2001 (20.07.2001) US

(71) Applicant (for all designated States except US): OY JU-  
VANTIA PHARMA LTD [FI/FI]; Lemminkäisenkatu 5,  
FIN-20520 Turku (FI).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JOUTSAMO,  
Topi [FI/FI]; Kohnmontie 10 B 37, FIN-20540 Turku (FI).

(81) Designated States (national): AE, AG, AL, AM, AT (uti-  
lity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (uti-  
lity model), DE, DK (utility model), DK, DM, DZ, EC, EE  
(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,  
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,  
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD,  
SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR,  
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

Published:

Puolikontie 6 C, FIN-20300 Turku (FI). HOFFRÉN,  
Anna-Marja [FI/FI]; Bilmarkinkatu 4 as 12, FIN-20100  
Turku (FI). WURSTER, Siegfried [DE/DE]; Matarakaari  
28 A 1, FIN-21500 Piikkiö (FI).

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS USEFUL FOR TREATMENT OR PREVENTION OF DISEASE MEDIATED BY AL-  
PHA-2B-ADRENOCEPTOR



(57) Abstract: The invention relates to compounds of formula (I) or a pharmaceutically acceptable salt thereof wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> are independently of each other H, a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen; X is H, a straight or branched alkyl chain with 1 to 4 carbon atoms, phenyl, -OH or =O; Z is H, acetyl, -CH<sub>2</sub>-Ph-O-CF<sub>3</sub> or CH<sub>2</sub>-Ph-CF<sub>3</sub>, Y is a ring structure optionally linked to formula (I) with an alkyl chain having one or two carbon

atoms. This invention further relates to the use of said compounds for the manufacture of a pharmaceutical preparation useful for the treatment or prevention of a disease mediated by the alpha-2B-adrenoceptor in a mammal.

WO 03/008387 A1

BEST AVAILABLE COPY

## **COMPOUNDS USEFUL FOR TREATMENT OR PREVENTION OF DISEASE MEDIATED BY ALPHA-2B-ADRENOCEPTOR**

The present invention relates to the use of selective alpha-2B-adrenoceptor antagonists for the manufacture of a pharmaceutical preparation useful for the treatment or prevention of diseases mediated by the alpha-2B-adrenoceptor in mammals.

### **BACKGROUND OF THE INVENTION**


The publications and other materials used herein to illuminate the background of the invention, and in particular, cases to provide additional details respecting the practice, are incorporated by reference.

It is known that alpha-2B-adrenoceptors mediate vascular contractions. Therefore, alpha-2B-antagonists are useful in the treatment or prevention of diseases involving vascular contraction. It has also been found that certain individuals have a genetic polymorphism in the alpha-2B-adrenoceptor gene. It has been observed that the alpha-2B-adrenoceptor protein in some subjects has a deletion of 3 glutamates from the glutamic acid repeat element of 12 glutamates (amino acids 297–309), in an acid stretch of 17 amino acids, located in the third intracellular loop of the receptor polypeptide (WO 01/29082; Heinonen et al., 1999).

### **OBJECTS AND SUMMARY OF THE INVENTION**

An object of this invention is to provide compounds useful for the treatment or prevention of a disease mediated by the alpha-2B-adrenoceptor in a mammal.

Thus this invention concerns a novel compound of formula (I)



or a pharmaceutically acceptable salt thereof.

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are independently of each other H, a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

5  $X$  is H, a straight or branched alkyl chain with 1 to 4 carbon atoms, phenyl or -OH;

$Z$  is H, acetyl,  $-\text{CH}_2\text{-Ph-O-CF}_3$  or  $-\text{CH}_2\text{-Ph-CF}_3$ , where Ph is phenyl;

$Y$  is a ring structure optionally linked to formula (I) with an alkyl chain having

10 a) phenyl optionally mono- or disubstituted and each substituent is independently selected from the group consisting of a halogen, a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, a halogen substituted methyl or methoxy group, a nitrile, an amide, amino, or a nitro group;


15 b) 2-benzimidazolyl, 2-imidazolyl, or 2- or 3-indolyl, wherein one N optionally has a substituent that is a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, or benzyl; and wherein the 2-benzimidazolyl, 2-imidazolyl, or 2- or 3-indolyl is optionally mono- or disubstituted and each substituent can independently be a straight or  
20 branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

(c) pyridinyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen; or

(d) naphthyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen.

The following previously known compounds are excluded: 4-[(1*H*-benzimidazol-2-ylmethyl)-amino]-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide (Kumar & Reddy, 1985), *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-methyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide (No 653716, ChemBridge Corporation, 16981 Via Tazon, Suite G, San Diego CA 92127) and *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide (No AE-848/34956037, SPECS and BioSPECS B. V., Fleminglaan 16, 2289 CP Rijswijk, The Netherlands) and *N*-(4-methyl-2-pyrimidinyl)-4-[(1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide (Farag & El-Mouafi & Khalifa, 1991).

THIS INVENTION FURTHER CONCERNS USE OF COMPOUNDS OF FORMULA (I)



15

or a pharmaceutically acceptable salt thereof for the manufacture of a pharmaceutical preparation useful for the treatment or prevention of a disease mediated by the alpha-2B-adrenoceptor in a mammal wherein

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are independently of each other H, a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

20

$X$  is H, a straight or branched alkyl chain with 1 to 4 carbon atoms, phenyl, -OH or =O;

$Z$  is H, acetyl, -CH<sub>2</sub>-Ph-O-CF<sub>3</sub> or -CH<sub>2</sub>-Ph-CF<sub>3</sub>, where Ph is phenyl;

Y is a ring structure optionally linked to formula (I) with an alkyl chain having one or two carbon atoms, wherein the ring structure is

5 a) phenyl optionally mono- or disubstituted and each substituent is independently selected from the group consisting of a halogen, a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, a halogen substituted methyl or methoxy group, an acetyl, a nitrile, an amide, amino, or a nitro group;

10 b) 2-benzimidazolyl, 2-imidazolyl, or 2- or 3-indolyl, wherein one N optionally has a substituent that is a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, or benzyl; and wherein the 2-benzimidazolyl, 2-imidazolyl, or 2- or 3-indolyl is optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

15 c) pyrimidinyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen; or


(d) naphthyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen.

20 The following compounds previously known to be selective alpha-2B-adrenoceptor antagonists are excluded: *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-methyl-1*H*-benzimidazol-2-ylmethyl)-amino] -benzenesulfonamide (No 653716, ChemBridge Corporation, 16981 Via Tazon, Suite G, San Diego CA 92127), *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-benzimidazol-2-ylmethyl)-amino] -benzenesulfonamide (No AE-848/34956037, SPECS and  
25 BioSPECS B. V., Fleminglaan 16, 2289 CP Rijswijk, The Netherlands) and *N*-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-4-ethoxy-benzamide (No AF-

399/36012031, SPECS and BioSPECS B. V., Fleminglaan 16, 2289 CP Rijswijk, The Netherlands).

## DETAILED DESCRIPTION OF THE INVENTION

Preferred compounds of the invention are compounds of formula (I)



5

as defined in the preceding summary or pharmaceutically acceptable salts thereof wherein  $R_1$  and  $R_3$  are methyl and  $R_2$ ,  $R_4$  and  $R_5$  are H.

In some preferable compounds  $X$  is H,  $Y$  is a phenyl optionally mono- or disubstituted with a straight or branched alkoxy group and  $Z$  is H. Compounds fulfilling all of the aforementioned characteristics and wherein said phenyl is substituted and said alkoxy substituent is methoxy are

4-(2,4-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methoxybenzylamino)-benzenesulfonamide,

4-(3,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide and *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methoxybenzylamino)-benzenesulfonamide.

In other preferred compounds  $X$  is H,  $Y$  is a phenyl optionally mono- or disubstituted with a straight or branched alkyl and/or a halogen and  $Z$  is H.

These comprise compounds such as 4-benzylamino-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methylbenzyl-

20




amino)-benzenesulfonamide, 4-(2,4-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methylbenzylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(4-methylbenzylamino)-benzenesulfonamide, 4-(2,5-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,6-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(4-bromobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide and 4-(2,6-dichlorobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide.

Further preferred compounds are *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-indol-3-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-isobutyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(1-phenylethylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[2-(2-methoxyphenyl)-ethylamino]-benzenesulfonamide.

According to one embodiment of the invention the compound is *N*-(4-methyl-2-pyrimidinyl)-4-[(1-methyl-1*H*-benzimidazol-2-yl)-methyl]amino]-benzenesulfonamide.

The invention also relates to the use of selective alpha-2B-adrenoceptor antagonists of formula (I)



as defined in the preceding summary, or a pharmaceutically acceptable salt thereof for the manufacture of a pharmaceutical preparation.

In many preferable compounds to be used  $R_1$  and  $R_3$  are typically methyl and  $R_2$ ,  $R_4$  and  $R_5$  are typically H.

- 5 In some compounds preferably used  $X$  is H,  $Y$  is a phenyl optionally mono- or disubstituted with a straight or branched alkoxy group and  $Z$  is H. Especially preferable for use are compounds in which said phenyl is substituted and said alkoxy substituent is methoxy. Such compound comprise 4-(2,4-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-  
 10 (4,6-dimethylpyrimidin-2-yl)-4-(3-methoxybenzylamino)-benzenesulfonamide, 4-(3,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,6-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(4-methoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,4-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methylbenzylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methylbenzylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(4-methylbenzylamino)-benzenesulfonamide, 4-(2,5-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,6-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(4-bromobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide and 4-(2,6-dichlorobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide.
- 15 In other compounds preferably used  $X$  is H,  $Y$  is a phenyl optionally mono- or disubstituted with a straight or branched alkyl and/or a halogen and  $Z$  is H. Such compound comprise 4-benzylamino-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methylbenzylamino)-benzenesulfonamide, 4-(2,4-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methylbenzylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(4-methylbenzylamino)-benzenesulfonamide, 4-(2,5-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,6-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(4-bromobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide and 4-(2,6-dichlorobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide.
- 20  
25



Further preferred compounds to be used comprise 4-[(1*H*-benzimidazol-2-ylmethyl)-amino]-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-indol-3-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-isobutyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(1-phenylethylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[2-(2-methoxyphenyl)-ethylamino]-benzenesulfonamide and *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(naphthalen-2-ylmethyl)-amino]-benzenesulfonamide.

Alpha-2B-adrenoceptor antagonists are useful in the treatment and/or prevention of many diseases.

Individuals having a deletion in the alpha-2B-adrenoceptor protein (WO 01/29082; Heinonen et al., 1999), particularly the deletion/deletion genotype (D/D genotype)

alpha-2B-adrenoceptor antagonists. These individuals have a deletion of 3 glutamates from the glutamic acid repeat element of 12 glutamates (amino acids 297–309), in an acid stretch of 17 amino acids, located in the third intracellular loop of the receptor polypeptide.

It has been found that in a population-based cohort of Finnish middle-aged men that subjects with a D/D genotype of the alpha-2B-adrenoceptor gene have a significantly elevated risk for acute myocardial infarction (AMI) in a five-year follow-up study. The risk for AMI was increased in subjects who had no previously diagnosed coronary heart disease (CHD) at the study outset. Therefore, it has been postulated that the D/D genotype is related to an impaired capacity to down-regulate alpha-2B-adrenoceptor function during sustained receptor activation. Therefore, alpha-2B-adrenoceptors are believed to be involved in the pathogenesis of a significant fraction of all cases of AMI, especially in subjects

with the D/D genotype, but also in I/D and I/I subjects (I means "insertion" and stands for the "normal" allele).

The alpha-2B-adrenoceptor antagonists as disclosed in this invention would be particularly useful in the treatment or prevention of coronary heart diseases. As

5 examples can be mentioned

a) Acute AMI

If alpha-2B-adrenoceptor dependent vasoconstriction is a causative factor in some cases of AMI, then antagonism of these receptors should restore coronary circulation and reduce the ischemic myocardial damage.

10 b) Unstable angina pectoris

An alpha-2B-adrenoceptor antagonist will relieve the vasoconstrictive component in the sustained ischemic episode, thus alleviating the symptoms

c) Prinzmetal's variant form of angina pectoris

15 Vasoconstriction is a key factor in the pathogenesis of Prinzmetal's angina, and an alpha-2B- adrenoceptor antagonist may resolve and prevent attacks.

d) Other forms of chronic angina pectoris and CHD

20 An alpha-2B-adrenoceptor antagonist will help to alleviate the vasoconstrictive component in all types of CHD, providing both symptomatic relief and protection from AMI. A general reduction in vascular tone will contribute to this by reducing venous return, cardiac workload and oxygen consumption (a nitrate-type effect; see below).

e) Prevention of restenosis after coronary angioplasty in cases where vasoconstriction plays a role in restenosis.

25 Furthermore, the alpha-2B-adrenoceptor antagonists as disclosed in this invention would be useful in the treatment or prevention of essential hypertension, especially

in subjects with increased sympathetic activity and a hyperdynamic circulatory system.

In the study mentioned above, the D/D variant of the alpha-2B-adrenoceptor gene was not clearly associated with blood pressure. The inventors believe that this was  
5 due to two main factors, 1) antihypertensive treatment, and 2) complex regulation of systemic blood pressure. In another study (Heinonen et al.), it was observed that the D/D genotype was associated with reduced basal metabolic rate and reduced heart rate. These associations probably reflect increased vascular resistance in these subjects.

10 In transgenic mice with targeted inactivation of the alpha-2B-adrenoceptor gene, intravenously administered alpha-2-adrenoceptor agonists fail to induce the characteristic blood pressure elevation, which is seen in normal animals and also in humans. In humans, doses of such drugs (Glick et al., 1996). The hypotensive  
15 effect of these drugs was markedly accentuated. This demonstrates that alpha-2B-adrenoceptors mediate vascular contraction. Thus, an antagonist should reduce blood pressure. This effect has not been seen with alpha-2B-non-selective alpha-2-adrenoceptor antagonists, because antagonism of alpha-2A-adrenoceptors increases sympathetic outflow, cardiac output and blood pressure. In mice with dysfunctional alpha-2A-adrenoceptors, alpha-2-adrenoceptor agonists caused an  
20 accentuated hypertensive response and no hypotension (MacMillan et al., 1996).

An alpha-2B-adrenoceptor antagonist is postulated to have favourable effects in hypertensive subjects through their effects on renal function, muscle blood flow, and also on vascular resistance in other vascular beds. The anti-AMI effect of such a drug will be an additional benefit, as hypertension is a significant risk factor for  
25 AMI. This protection is due to three factors: 1) a reduction in systemic blood pressure, 2) decreased risk of coronary vasoconstriction, and 3) a nitrate-like effect on venous return, myocardial workload and oxygen consumption.

Moreover, the alpha-2B-adrenoceptor antagonists as disclosed in this invention would be useful in the treatment or prevention of other vascular diseases. Specifically, benefits can be expected in the treatment or prevention of

- vasoconstriction and hypoxic brain damage subsequent to subarachnoid
- 5 haemorrhage,
- migraine,
- Raynaud's disease and cold intolerance,
- pre-eclampsia,
- male erectile dysfunction, and
- 10 - obesity and the metabolic syndrome.

The last mentioned effect is due to the fact that reduced muscle blood flow and reduced basal metabolic rate contribute to the development of obesity and hypertension. An alpha-2B-adrenoceptor antagonist will, by increasing the muscle

15 favourable direction.

The alpha-2B-adrenoceptor antagonists disclosed in this invention are also useful in anaesthesia and analgesia to potentiate the clinical efficacy of alpha-2-adrenoceptor agonists, which are not selective for the alpha-2B-adrenoceptor subtype. By blocking the vasoconstriction induced by these agonists, a

20 simultaneously administered alpha-2B-adrenoceptor antagonist will allow the use of larger doses of said agonists, up to anaesthetic dose levels which have not previously been possible in man, only in veterinary anaesthetic practice.

## EXPERIMENTAL SECTION

**Human alpha-2-adrenoceptor binding affinity**

The affinity of test compounds for the three human  $\alpha_2$ -adrenoceptor subtypes ( $\alpha_{2A}$ ,  $\alpha_{2B}$  and  $\alpha_{2C}$ ) was determined in competition binding assays with  $^3\text{H}$ -rauwolscine.

5 The biological material for these experiments consisted of membranes from Shionogi S115 cells stably transfected with any of the three human  $\alpha_2$  subtypes (Marjamäki et al. 1992). Membrane (5–10  $\mu\text{g}$  of total protein per sample) and 1–2 nM  $^3\text{H}$ -rauwolscine (specific activity 78 Ci/mmol) were incubated in 50 mM  $\text{KH}_2\text{PO}_4$ , pH 7.5 with 6 concentrations of the compounds. Each concentration was

10 run in duplicate. Non-specific binding was defined by 100  $\mu\text{M}$  oxymetazoline and corresponded to 5–15% of total binding. After 30 min at room temperature, incubations were terminated by rapid vacuum filtration through GF/B glass fiber filter and three times washes with ice cold incubation buffer. The filters were then

dried, impregnated with scintillate and their radioactivity was measured by

15 scintillation counting. The analysis of the experiments was carried out by non-linear least square curve fitting. Experimentally determined  $\text{IC}_{50}$  values were converted to  $\text{K}_i$ 's by making use of the Cheng-Prusoff equation (Cheng and Prusoff, 1973). Experiments were repeated a minimum of three times.

**Table 1:** Human  $\alpha_2$ -adrenoceptor subtypes binding affinities. Data is presented as

20  $\text{K}_i$ 's in nM (Mean  $\pm$  SEM).

Compound	alpha-2A	alpha-2B	alpha-2C
A	>13000	160 $\pm$ 20	>30000
B	>4500	34 $\pm$ 2	>10000
C	2000 $\pm$ 400	10 $\pm$ 2	>10000
D	>10000	440 $\pm$ 70	>10000
E	>5100	20 $\pm$ 4	>10000

F	>4300	$43 \pm 7$	>10000
G	$2200 \pm 600$	$32 \pm 5$	>10000
H	>30000	$8000 \pm 500$	>30000

Results expressed in the form of ">" means that no numerical values for  $K_i$ 's could be established due to lack of displacement or due to incomplete competition curves. However, the experimental data indicated that, at a minimum, the  $K_i$ 's must be larger than the numbers given.

#### 5 Antagonist activity on human alpha-2-adrenoceptor subtypes

Antagonist potencies were determined as the ability of test compounds to competitively inhibit epinephrine-stimulated  $^{35}\text{S}$ -GTP $\gamma$ S binding to G proteins (Tian et al., 1993; Wieland and Jakobs, 1994; Jasper et al., 1998) in membranes of CHO cells stably transfected with one of the three human  $\alpha_2$ -subtypes (Pohjanpelto et al., 1997; Marjamäki et al., 1998). Membranes (2–6  $\mu\text{g}$  of protein per sample) and 12 concentrations of test compound were preincubated for 30 min with a fixed concentration of epinephrine (5  $\mu\text{M}$  for  $\alpha_{2A}$ , 15  $\mu\text{M}$  for  $\alpha_{2B}$ , 5  $\mu\text{M}$  for  $\alpha_{2C}$ ) in 50 mM Tris, 5 mM  $\text{MgCl}_2$ , 150 mM NaCl, 1 mM DTT, 1 mM EDTA, 10  $\mu\text{M}$  GDP, 30  $\mu\text{M}$  ascorbic acid, pH 7.4 at room temperature. Binding of radiolabel was started by the addition of trace amounts of  $^{35}\text{S}$ -GTP $\gamma$ S (0.08–0.15 nM, specific activity 1250 Ci/mmol) to the incubation mixture. After an additional 60 min at room temperature, the incubation was terminated by rapid vacuum filtration through glass fibre filter. Filters were washed three times with 5 ml ice cold wash buffer (20 mM Tris, 5 mM  $\text{MgCl}_2$ , 1 mM EDTA pH 7.4 at room temperature), dried and counted for radioactivity in a scintillation counter. Analysis of experiments was carried out by non-linear least square fitting. Results are based on a minimum of three experiments.



**Table 2:** Antagonist effect on human  $\alpha_2$ -adrenoceptor subtypes. Data is presented as KB's in nM (Mean  $\pm$  SEM).

Compound	alpha-2A	alpha-2B	alpha-2C
B	14500 $\pm$ 3600	75 $\pm$ 9	5700 $\pm$ 700
C	5400 $\pm$ 1400	17 $\pm$ 5	6300 $\pm$ 1400
E	7900 $\pm$ 3100	29 $\pm$ 5	7300 $\pm$ 1100
F	8700 $\pm$ 1100	240 $\pm$ 60	12000 $\pm$ 2000
G	3200 $\pm$ 500	86 $\pm$ 64	4700 $\pm$ 1800

For the purpose of the invention, the alpha-2B-adrenoceptor antagonist or its pharmaceutically acceptable salt can be administered by various routes. The suitable administration forms include, for example, oral formulations; parenteral injections including intravenous, intramuscular, intradermal and subcutaneous injections; transdermal or rectal administration forms. The required dosage of the compounds of the alpha-2B-adrenoceptor antagonist will vary with the particular condition being treated, the severity of the condition, the duration of the treatment, the administration route and the specific compound being employed. The suitable dose varies in the range 5  $\mu$ g to 100 mg per kg body weight and day for an adult person.

## EXAMPLES

**Example 1**

*N*-(4,6-Dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-benzimidazol-2-ylmethyl)-amino]  
-benzenesulfonamide

**5    *Step I: Alkylation of 2-hydroxymethylbenzimidazole***

99.2 mg (0.67 mmol) 2-hydroxymethylbenzimidazole was dissolved in 3 ml methanol. Potassium carbonate (103.1 g, 0.75 mmol) and diethylsulfate (442 µl 3.38 mmol) were added to the reaction mixture. Solution was stirred and refluxed overnight. The reaction mixture was then evaporated to dryness and purified on  
10    silica using gradient elution (chloroform to 5% methanol in chloroform) to obtain

***Step II: Chlorination of 1-ethyl-2-hydroxymethylbenzimidazole***

20 mg (0.11 mmol) 1-ethyl-2-hydroxymethylbenzimidazole was dissolved in 2 ml dichloromethane. Thionyl chloride (24 µl, 0.33 mmol) was diluted 20 times with  
15    dichloromethane and the solution was added to the reaction mixture. Reaction mixture was stirred at room temperature for two hours, evaporated to dryness and washed with water to yield 1-ethyl-2-chloromethylbenzimidazole as pale yellow crystals, 36 mg (95%).

***Step III: Coupling reaction between 1-ethyl-2-chloromethylbenzimidazole and  
20    sulfamethazine***

32.4 mg (0.12 mmol) sulfamethazine and 36 mg (0.18 mmol) 1-ethyl-2-chloromethylbenzimidazole were dissolved in 4 ml methanol. 64 µl (0.44 mmol) triethylamine and catalytic amount of sodium iodide were added to the reaction

mixture. Solution was stirred and refluxed overnight. The reaction mixture was then evaporated to dryness, and purified on silica using gradient elution (chloroform to 5% methanol in chloroform) to provide white crystals of the title compound, 10 mg (20%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 7.68 (2H, m), 7.60 (1H, m), 7.52 (1H, m), 7.22 (1H, m), 7.17 (1H, m), 7.00 (1H, br, s), 6.78 (3H, m), 6.60 (1H, br, s), 4.60 (2H, m), 4.30 (2H, q, 7.2 Hz), 2.18 (6H, s), 1.28 (3H, t, 7.2 Hz); MS (ESI<sup>+</sup>): m/z 437 (M + H)<sup>+</sup>.

### Example 2

#### 10 4-[(1H-Benzimidazol-2-ylmethyl)-amino]-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

Following the procedure outlined in Step III of example 1, but without triethylamine and substituting 1-ethyl-2-chloromethylbenzimidazole by 2-chloromethylbenzimidazole afforded the title compound with the yield of 51%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 7.70 (2H, m), 7.48 (2H, br, m), 7.13 (2H, m), 7.11 (1H, t, 5.8 Hz), 6.70 (3H, m), 4.52 (2H, d, 5.8 Hz), 2.21 (6H, s); MS (ESI<sup>+</sup>): m/z 409 (M + H)<sup>+</sup>.

### Example 3

20 N-(4,6-Dimethylpyrimidin-2-yl)-4-[(pyridin-4-ylmethyl)-amino]-benzenesulfonamide

Following the procedure outlined in Step III of example 1 without triethylamine and substituting 1-ethyl-2-chloromethylbenzimidazole by 4-picolylchloride hydrochloride afforded the title compound with the yield of 54%. MS (ESI<sup>+</sup>): m/z 392 (M + Na)<sup>+</sup>, 370 (M + H)<sup>+</sup>.

**Example 4**

*N*-(4,6-Dimethylpyrimidin-2-yl)-4-[(1-isobutyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide (Compound A)

Following the procedure outlined in example 1, but substituting in step I ethyl  
5 bromide for isobutyl iodide, afforded the title compound with stepwise yields of  
15%, 95% and 15%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 7.76 (2H, m), 7.70 (1H, m),  
7.60 (1H, m), 7.27 (1H, m), 7.22 (1H, m), 7.15 (1H, br, t, 5.3 Hz), 6.86 (2H, m),  
6.77 (1H, s), 4.65 (2H, d, 5.3 Hz), 4.13 (2H, d, 7.5 Hz), 2.27 (6H, s), 2.25 (1H, m),  
0.91 (6H, d, 6.7 Hz); MS (ESI<sup>+</sup>): m/z 487 (M + Na)<sup>+</sup>, 465 (M + H)<sup>+</sup>.

**10 Example 5**

yl)- benzenesulfonamide

Following the procedure outlined in example 1, but substituting in step I ethyl  
bromide for benzyl bromide, afforded the title compound with stepwise yields of  
15 23%, 90% and 18%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 7.81 (2H, m), 7.52 (1H, m),  
7.45 (1H, m), 7.30 (5H, m), 7.16 (2H, m), 6.75 (1H, s), 6.54 (2H, m), 6.02 (1H, br,  
s), 5.66 (2H, s), 5.63 (2H, s), 2.21 (6H, s); MS (ESI<sup>+</sup>): m/z 521 (M + Na)<sup>+</sup>, 499 (M  
+ H)<sup>+</sup>.

**Example 6**

20 4-[(1-Ethyl-1*H*-benzimidazol-2-ylmethyl)-amino]-*N*-(5-methoxypyrimidin-2-yl)-benzenesulfonamide

Following the procedure outlined in example 1 step III, but substituting  
sulfamethazine by 5-methoxysulfadiazine, afforded the title compound with the

yield of 8%. Dimethylformamide was used as a solvent and additional silica gel chromatography purification with 2:1 petrol ether:ethylacetate was needed. MS (ESI<sup>+</sup>): m/z 461 (M + Na)<sup>+</sup>, 439 (M + H)<sup>+</sup>.

### Example 7

- 5 4-[(1H-Benzimidazol-2-ylmethyl)-amino]-N-(pyrimidin-2-yl)-benzenesulfonamide  
628 mg (3.8 mmol) sulfadiazine and 728 mg (3.0 mmol) 2-chloromethylbenzimidazole were dissolved in 10 ml 1 M NaOH. Solution was stirred and refluxed for four hours. Reaction mixture was neutralised with addition of 1 M acetic acid until product precipitated. Crystals were filtered and purified on silica  
10 using gradient elution (chloroform to 5% methanol in chloroform) to obtain the title compound as white crystals with 38% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 6.98 (1H, m), 6.72 (2H, m), 4.53 (2H, 5.7 Hz); MS (ESI<sup>+</sup>): m/z 381 (M + H)<sup>+</sup>.

### Example 8

- 15 N-(1H-Benzimidazol-2-ylmethyl)-N-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-acetamide  
18 mg (0.044 mmol) 4-[(1H-benzimidazol-2-ylmethyl)-amino]-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide was dissolved in 2 ml of 15% pyridine in dichloromethane. Acetyl chloride (31 μl, 0.44 mmol) was diluted with 1 ml  
20 dichloromethane and solution was added to the reaction mixture. After three hours reaction mixture was washed with acidic water and organic layer was evaporated to dryness. Crystals were purified on silica using gradient elution (chloroform to 5% methanol in chloroform) to obtain white crystals with 30% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 7.97 (2H, m), 7.60 (2H, m), 7.48 (2H, m), 7.14 (2H, m),

6.68 (1H, s), 5.08 (2H, s), 2.18 (6H, s), 1.93 (3H, s); MS (ESI<sup>+</sup>): m/z 473 (M + Na)<sup>+</sup>, 451 (M + H)<sup>+</sup>.

### Example 9

5 *N*-(1-Acetyl-1*H*-benzimidazol-2-ylmethyl)-*N*-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-acetamide

Title compound was purified from the reaction mixture produced according to example 8 with a yield of 14%. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 8.15 (2H, m), 7.80 (2H, m), 7.55 (2H, m), 7.48 (1H, s), 7.21 (3H, m), 5.17 (2H, s), 2.54 (6H, s), 2.03 (3H, s), 1.83 (3H, s); MS (ESI<sup>+</sup>): m/z 493 (M + H)<sup>+</sup>.

10 **Example 10**

4-[(1-Acetyl-1*H*-benzimidazol-2-ylmethyl)-amino]-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

Following the procedure of example 8, but instead of 15% pyridine in dichloromethane only few drops of pyridine in dichloromethane were used as a  
15 solvent. Method afforded the title compound with a yield similar to that for *N*-(1*H*-benzimidazol-2-ylmethyl)-*N*-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-acetamide. MS (ESI<sup>+</sup>): m/z 451 (M + H)<sup>+</sup>.

### Example 11

4-Benzylamino-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

20 100 mg (0.36 mmol) sulfamethazine and 70.8 μl (0.60 mmol) benzyl bromide were dissolved in 4 ml methanol. Caesium carbonate (113.4 mg, 0.35 mmol) was added and solution was refluxed overnight with stirring. The reaction mixture was then



evaporated to dryness, and purified on silica using gradient elution (chloroform to 2% methanol in chloroform) to obtain white crystals in a yield similar to that described in example 1 for step III. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.93 (2H, m), 7.31 (5H, m), 6.58 (3H, m), 4.36 (2H, s), 2.34 (6H, s); MS (ESI<sup>+</sup>): m/z 369 (M + H)<sup>+</sup>.

### Example 12

#### 4-(4-Bromobenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

Following the procedure outlined in example 11, but substituting benzyl bromide by 4-bromobenzyl bromide, afforded the title compound with a yield of 8%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.91 (2H, m), 7.45 (2H, m), 7.19 (2H, m), 6.59 (1H, s), 6.55 (2H, m), 4.32 (2H, s), 2.34 (6H, s); MS (ESI<sup>+</sup>): m/z 469 (M + Na)<sup>+</sup>, 447 (M + H)<sup>+</sup>.

### Example 13

#### N-(4,6-Dimethylpyrimidin-2-yl)-4-(2-methylbenzylamino)-benzenesulfonamide

#### 15 (Compound B)

Following the procedure outlined in example 11, but substituting benzyl bromide by 2-methylbenzyl bromide, afforded the title compound in a yield similar to that described in example 1 for step III. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.95 (2H, m), 7.22 (4H, m), 6.58 (3H, m), 4.30 (2H, s), 2.35 (3H, s), 2.34 (6H, s); MS (ESI<sup>+</sup>): m/z 405 (M + Na)<sup>+</sup>.

**Example 14***N*-(4,6-Dimethylpyrimidin-2-yl)-4-(4-methylbenzylamino)-benzenesulfonamide

Following the procedure outlined in example 11, but substituting benzyl bromide by 4-methylbenzyl bromide, afforded the title compound in a yield similar to that  
5 described in example 1 for step III. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.92 (2H, m), 7.20 (2H, m), 7.15 (2H, m), 6.58 (3H, m), 4.31 (2H, s), 2.34 (3H, s), 2.33 (6H, s); MS (ESI<sup>+</sup>): m/z 405 (M + Na)<sup>+</sup>, 383 (M + H)<sup>+</sup>.

**Example 15***N*-(4,6-Dimethylpyrimidin-2-yl)-4-(1-phenylethylamino)-benzenesulfonamide

10 Following the procedure outlined in example 11, but substituting benzyl bromide by (1-bromoethyl)-benzene, afforded the title compound with a yield of 12%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.84 (2H, m), 7.31 (5H, m), 6.57 (1H, s), 6.46 (2H, m), 4.53 (1H, q, 6.7 Hz), 2.30 (6H, s), 1.54 (3H, d, 6.7 Hz); MS (ESI<sup>+</sup>): m/z 405 (M + Na)<sup>+</sup>, 383 (M + H)<sup>+</sup>.

**15 Example 16***N*-(4,6-Dimethylpyrimidin-2-yl)-4-(2-methoxybenzylamino)-benzenesulfonamide

Following the procedure outlined in example 11, but substituting benzyl bromide by 2-methoxybenzyl bromide, afforded the title compound in a yield similar to that described in example 1 for step III. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.92 (2H, m),  
20 7.25 (2H, m), 6.90 (2H, m), 6.59 (3H, m), 4.36 (2H, s), 3.86 (3H, s), 2.32 (6H, s); MS (ESI<sup>+</sup>): m/z 421 (M + Na)<sup>+</sup>, 399 (M + H)<sup>+</sup>.

**Example 17****4-(2,4-Dimethylbenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide**

Following the procedure outlined in example 11, but substituting benzyl bromide  
5 by 2,4-dimethylbenzyl bromide, afforded the title compound with a yield of 23%.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.94 (2H, m), 7.14 (1H, m), 7.03 (1H, s), 6.98 (1H, m), 6.58 (3H, m), 4.26 (2H, s), 2.34 (6H, s), 2.31 (3H, s), 2.30 (3H, s); MS (ESI<sup>+</sup>): m/z 419 (M + Na)<sup>+</sup>, 397 (M + H)<sup>+</sup>.

**Example 18****N-(4,6-Dimethylpyrimidin-2-yl)-4-(3-methylbenzylamino)-benzenesulfonamide**

Following the procedure outlined in example 11, but substituting benzyl bromide  
by 3-methylbenzyl bromide, afforded the title compound with a yield of 10%. <sup>1</sup>H  
NMR (CDCl<sub>3</sub>, 500 MHz): 7.93 (2H, m), 7.23 (1H, m), 7.11 (3H, m), 6.62 (1H, s),  
6.58 (2H, m), 4.31 (2H, s), 2.36 (6H, s), 2.34 (3H, s); MS (ESI<sup>+</sup>): m/z 405 (M +  
15 Na)<sup>+</sup>, 383 (M + H)<sup>+</sup>.

**Example 19****4-(2,6-Dichlorobenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide  
(Compound C)**

Following the procedure outlined in example 11, but substituting benzyl bromide  
20 by 2,6-dichlorobenzyl bromide, afforded the title compound with the yield of 10%.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.96 (2H, m), 7.34 (2H, m), 7.20 (1H, m), 6.71 (2H, m), 6.59 (1H, s), 4.63 (2H, s), 2.34 (6H, s); MS (ESI<sup>+</sup>): 437 (M + H)<sup>+</sup>.

**Example 20**

*N*-(4,6-Dimethylpyrimidin-2-yl)-4-[(naphthalen-2-ylmethyl)-amino]-benzenesulfonamide (Compound D)

Following the procedure outlined in example 11, but substituting benzyl bromide  
5 by 2-bromomethylnaphtalene, afforded the title compound with the yield of 20%.  
<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 500 MHz): 7.85 (4H, m), 7.68 (2H, m), 7.47 (3H, m), 7.19  
(1H, t, 5.8 Hz), 6.70 (1H, s), 6.66 (2H, m), 4.49 (2H, d, 5.8 Hz), 2.21 (6H, s); MS  
(ESI<sup>+</sup>): m/z 441 (M + H)<sup>+</sup>.

**Example 21**

10 *N*-(4,6-Dimethylpyrimidin-2-yl)-4-(3-methoxybenzylamino)-benzenesulfonamide

Following the procedure outlined in example 11, but substituting benzyl bromide  
by 3-methoxybenzyl bromide, afforded the title compound with the yield of 28%.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.93 (2H, m), 6.86 (3H, m), 6.58 (3H, m), 4.33 (2H,  
15 s), 3.78 (3H, s), 2.33 (6H, s); MS (ESI<sup>+</sup>): m/z 421 (M + Na)<sup>+</sup>, 399 (M + H)<sup>+</sup>.

**Example 22**

*N*-(4,6-Dimethylpyrimidin-2-yl)-4-(4-nitrobenzylamino)-benzenesulfonamide

Following the procedure outlined in example 11, but substituting benzyl bromide  
by 4-nitrobenzyl bromide, afforded the title compound with a yield lower than 1%.  
20 MS (ESI<sup>+</sup>): m/z 414 (M + H)<sup>+</sup>.

**Example 23**

*N*-(4,6-Dimethylpyrimidin-2-yl)-4-(4-trifluoromethylbenzylamino)-  
benzenesulfonamide

Following the procedure outlined in example 11, but substituting benzyl bromide  
5 by 4-trifluoromethylbenzyl bromide, afforded the title compound in a yield similar  
to that described in example 1 for step III. MS (ESI<sup>+</sup>): m/z 459 (M + Na)<sup>+</sup>, 437 (M  
+ H)<sup>+</sup>.

**Example 24**

4-[Bis-(4-trifluoromethylbenzyl)-amino]-*N*-(4,6-dimethylpyrimidin-2-yl)-  
10 benzenesulfonamide

Title compound was purified from the reaction mixture produced according to  
example 23 with the yield of 20%. MS (ESI<sup>+</sup>): 617 m/z (M + Na)<sup>+</sup>, 595 (M + H)<sup>+</sup>.

**Example 25**

*N*-(4,6-Dimethylpyrimidin-2-yl)-4-(4-trifluoromethoxybenzylamino)-  
15 benzenesulfonamide

Following the procedure outlined in example 11, but substituting benzyl bromide  
by 4-trifluoromethoxybenzyl bromide, afforded the title compound with a yield of  
25%. MS (ESI<sup>+</sup>): m/z 475 (M + Na)<sup>+</sup>, 453 (M + H)<sup>+</sup>.

**Example 26**

4-[Bis-(4-trifluoromethoxybenzyl)-amino]-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

Title compound was purified from the reaction mixture produced according to  
5 example 25 with a yield of 15%. MS (ESI<sup>+</sup>): m/z 649 (M + Na)<sup>+</sup>, 627 (M + H)<sup>+</sup>.

**Example 27**

4-(2,5-Dimethylbenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

Following the procedure outlined in example 11 without caesium carbonate and  
10 substituting benzyl bromide by 2,5-dimethylbenzyl bromide afforded the title  
compound with a yield of 35%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.95 (2H, m), 7.07  
(3H, m), 6.59 (3H, m), 4.25 (2H, s), 2.33 (6H, s), 2.30 (3H, s), 2.28 (3H, s); MS  
(ESI<sup>+</sup>): m/z 419 (M + Na)<sup>+</sup>.

**Example 28**

15 4-(2,6-Dimethylbenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

Following the procedure outlined in example 11 without caesium carbonate and  
substituting benzyl bromide by 2,6-dimethylbenzyl bromide afforded the title  
compound with a yield of 25%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.99 (2H, m), 7.08  
20 (3H, m), 6.62 (3H, m), 4.26 (2H, s), 2.37 (12H, m); MS (ESI<sup>+</sup>): m/z 419 (M +  
Na)<sup>+</sup>.



**Example 29****4-(3,5-Dimethoxybenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide**

Following the procedure outlined in example 11 without caesium carbonate and substituting benzyl bromide by 3,5-dimethoxybenzyl chloride afforded the title compound with a yield of 15%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.92 (2H, m), 6.60 (1H, s), 6.58 (2H, m), 6.46 (2H, m), 6.38 (1H, m), 4.30 (2H, s), 3.76 (6H, s), 2.33 (6H, s); MS (ESI<sup>+</sup>): m/z 429 (M + H)<sup>+</sup>.

**Example 30****4-(2,5-Dimethoxybenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide**

Following the procedure outlined in example 11 without caesium carbonate and substituting benzyl bromide by 2,5-methoxybenzyl chloride afforded the title compound with the yield of 13%. Reaction time was three days. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.91 (2H, m), 6.79 (3H, m), 6.58 (3H, m), 4.33 (2H, s), 3.81 (3H, s), 3.71 (3H, s), 2.33 (6H, s); MS (ESI<sup>+</sup>): m/z 429 (M + H)<sup>+</sup>.

**Example 31****2,6-Dichloro-N-[4-(4,6-dimethylpyrimidin-2-yl)sulfamoyl]-phenyl]-benzamide**

Following the procedure outlined in example 11, but substituting benzyl bromide by 2,6-dichlorobenzoyl chloride, afforded the title compound with almost quantitative yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 8.11 (2H, m), 7.80 (2H, m), 7.33 (3H, m), 6.59 (1H, s), 2.34 (6H, s); MS (ESI<sup>+</sup>): m/z 473 (M + H)<sup>+</sup>, 451 (M + H)<sup>+</sup>.

**Example 32****4-(2-Cyanobenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide**

Following the procedure outlined in example 11, but substituting benzyl bromide by  $\alpha$ -bromo-*o*-tolunitrile, afforded the title compound with a yield of 21%. <sup>1</sup>H  
5 NMR (DMSO-d<sub>6</sub>, 500 MHz): 7.83 (1H, m), 7.71 (2H, m), 7.65 (1H, m), 7.47 (2H, m), 7.15 (1H, br, t, 5.8 Hz), 6.73 (1H, m), 6.63 (2H, m), 4.50 (2H, d, 5.8 Hz), 2.24 (6H, s); MS (ESI<sup>+</sup>): m/z 416 (M + Na)<sup>+</sup>, 394 (M + H)<sup>+</sup>.

**Example 33****4-(2,4-Dimethoxybenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-  
10 benzenesulfonamide**

62 mg (0.37 mmol) 2,4-dimethoxybenzylaldehyde and 100.9 mg (0.50 mmol) sulfamethazine were dissolved in 4 ml of 1,2-dichloroethane. Acetic acid (168  $\mu$ l, 2.8 mmol) was added to the reaction mixture and solution was stirred with reflux overnight. Sodium triacetoxyborohydride (162.9 mg, 0.77 mmol) was dissolved to  
15 the reaction solution and refluxing was continued for three hours. The reaction mixture was then evaporated to dryness, and purified on silica using gradient elution (chloroform to 2% methanol in chloroform) and 2:1 petrol ether:ethylacetate to obtain the title compound as white crystals with a yield of 10%. . <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.91 (2H, m), 7.13 (1H, m), 6.58 (3H, m),  
20 6.47 (1H, m), 6.42 (1H, m), 4.53 (1H, t, 5.0 Hz), 4.30 (2H, d, 5.0 Hz), 3.82 (3H, s), 3.79 (3H, s), 2.33 (6H, s); MS (ESI<sup>+</sup>): m/z 429 (M + H)<sup>+</sup>.

**Example 34**

*N*-(4,6-Dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-indol-3-ylmethyl)-amino]-benzenesulfonamide (Compound F)

1-ethylindole-3-carboxaldehyde was prepared from alkylation reaction of indole-3-carboxaldehyde. Indole-3-carboxaldehyde (900 mg, 6.2 mmol) was dissolved in 5 ml dimethylformamide, ethyl bromide (918  $\mu$ l, 12 mmol) and sodium hydride (282.8 mg, 12 mmol) were added to the reaction mixture. Solution was stirred and refluxed for three hours. The reaction mixture was then evaporated to dryness and washed with water. Pale brown crystals were obtained with 80% yield.

Following the procedure outlined in example 33, but substituting 2,4-dimethoxybenzylaldehyde by 1-ethylindole-3-carboxaldehyde, afforded the compound with a <sup>1</sup>H NMR spectrum similar to that of *N*-(4-(2,4-dimethoxybenzylamino))-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide. Instead of sodium triacetoxyborohydride sodium borohydride was used to reduce the imine intermediate. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.88 (2H, m), 7.52 (1H, m), 7.29 (1H, m), 7.18 (1H, m), 7.05 (1H, m), 7.03 (1H, s), 6.56 (2H, m), 6.52 (1H, s), 4.41 (2H, d, 4.7 Hz), 4.33 (1H, t, 4.7 Hz), 4.08 (2H, q, 7.3 Hz), 2.27 (6H, s), 1.38 (3H, t, 7.3 Hz); MS (ESI<sup>+</sup>): *m/z* 458 (M + Na)<sup>+</sup>.

**Example 35**

*N*-[4-(4,6-Dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-2-(2-methoxyphenyl)-acetamide

2-Methoxyphenylacetic acid (33 mg, 0.20 mmol) and sulfamethazine (55 mg, 0.20 mmol) were dissolved in 4 ml of chloroform. Triethylamine (70  $\mu$ l, 0.55 mmol) and diisopropylcarbodiimide (50  $\mu$ l, 0.20 mmol) were added to the reaction mixture. Solution was stirred and refluxed overnight. The reaction mixture was

then evaporated to dryness, washed with water and purified on silica using gradient elution (chloroform to 2% methanol in chloroform) to obtain white crystals with the yield of 40%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 8.04 (2H, m), 7.54 (2H, m), 7.30 (2H, m), 6.98 (2H, m), 6.60 (1H, s), 3.95 (3H, s), 3.72 (2H, s), 2.33 (6H, s); MS (ESI<sup>+</sup>): m/z 449 (M + Na)<sup>+</sup>, 427 (M + H)<sup>+</sup>.

### Example 36

#### 2-Acetyl-N-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-benzamide

Following the procedure outlined in example 35, but substituting 2-methoxyphenylacetic acid by 2-acetylbenzoic acid, afforded the title compound with a yield of 11%. MS (ESI<sup>+</sup>): m/z 447 (M + Na)<sup>+</sup>, 425 (M + H)<sup>+</sup>.

### Example 37

#### 1-Methyl-1H-indole-2-carboxylic acid [4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-amide

Following the procedure outlined in example 35, but substituting 2-methoxyphenylacetic acid by 1-methylindole-2-carboxylic acid and using dimethylformamide as a solvent afforded the title compound with a yield similar to that of N-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-2-(2-methoxyphenyl)-acetamide. MS (ESI<sup>+</sup>): m/z 458 (M + Na)<sup>+</sup>.

**Example 38**2-(3,5-Dimethoxyphenyl)-N-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-acetamide

Following the procedure outlined in example 35, but substituting 2-methoxyphenylacetic acid by 3,5-dimethoxyphenylacetic acid, afforded the title compound with a yield of 54%. MS (ESI<sup>+</sup>): m/z 457 (M + H)<sup>+</sup>.

**Example 39**N-(4,6-Dimethylpyrimidin-2-yl)-4-[2-(2-methoxyphenyl)-ethylamino]-benzenesulfonamide (Compound G)

33 mg (0.08 mmol) N-[4-(4,6-dimethylpyrimidin-2-ylsulfomoyl)-phenyl]-2-(2-methoxyphenyl)-acetamide was dissolved in 2 ml tetrahydrofuran. To the solution, a borane tetrahydrofuran complex was added to the reaction mixture under nitrogen atmosphere and solution was stirred overnight with reflux. Reaction was quenched with 6 M HCl and the reaction mixture was neutralised with 1 M NaOH. Product was extracted with chloroform and purified on silica using gradient elution (chloroform to 2% methanol in chloroform) to obtain white crystals with a yield of 20%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.92 (2H, m), 7.22 (1H, m), 7.12 (1H, m), 6.89 (2H, m), 6.60 (1H, br, s), 6.54 (2H, m), 3.85 (3H, s), 3.38 (2H, t, 6.9 Hz), 2.93 (2H, t, 6.9 Hz), 2.33 (6H, s); MS (ESI<sup>+</sup>): m/z 435 (M + Na)<sup>+</sup>, 413 (M + H)<sup>+</sup>.

**Example 40**4-[2-(3,5-Dimethoxyphenyl)-ethylamino]-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

Following the procedure outlined in example 39, but substituting *N*-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-2-(2-methoxyphenyl)-acetamide by 2-(3,5-dimethoxyphenyl)-*N*-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-acetamide, afforded the title compound with a yield of 11%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): 7.92 (2H, m), 6.58 (1H, s), 6.54 (2H, m), 6.34 (3H, s), 4.20 (1H, t, 5.8 Hz), 3.76 (6H, s), 3.42 (2H, m), 2.85 (2H, t, 6.9 Hz), 2.34 (6H, s); MS (ESI<sup>+</sup>): m/z 465 (M + Na)<sup>+</sup>, 443 (M + H)<sup>+</sup>.

**Example 41**4-(Benzhydrylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide (Compound H)

Sulfamethazine (105 mg, 0.38 mmol) and diphenylchloromethane (77 μl, 0.38 mmol) were dissolved in 3 ml pyridine. Solution was stirred and refluxed overnight. The reaction mixture was evaporated to dryness and dissolved to 1 M NaOH. Product was precipitated with 1 M acetic acid. Precipitation cycle was repeated for three times to give white crystals of the title compound with a yield of 8%. MS (ESI<sup>+</sup>): m/z 467(M + Na)<sup>+</sup>, 445 (M + H)<sup>+</sup>.

**Example 42**4-(4-Aminobenzylamino)-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide

18 mg (0.04 mmol) of *N*-(4,6-dimethylpyrimidin-2-yl)-4-(4-nitrobenzylamino)-benzenesulfonamide was dissolved in 1 ml of tetrahydrofurane. Hydrazinium



hydrate (50  $\mu$ l, 1.5 mmol) and catalytic amount of palladium on charcoal were added to the reaction mixture. Solution was stirred overnight. The reaction mixture was evaporated to dryness and purified on silica using gradient elution (chloroform to 4% methanol in chloroform) to obtain with a yield of 59%. MS  
5 (ESI<sup>+</sup>): m/z 406 (M + H)<sup>+</sup>, 384 (M + H)<sup>+</sup>.

#### Example 43

4-{[4-(4,6-Dimethylpyrimidin-2-ylsulfamoyl)-phenylamino]-methyl}-benzamide  
4-{[4-(4,6-Dimethylpyrimidin-2-ylsulfamoyl)-phenylamino]-methyl}-benzoic acid  
methyl ester was prepared like described on example 11 and was treated with 25%  
10 ammonia to obtain the title compound with a yield lower than 1%. <sup>1</sup>H NMR  
(CDCl<sub>3</sub>, 500 MHz): 7.81 (2H, m), 7.67 (2H, m), 7.37 (2H, m), 7.12 (1H, br, t, 6.0  
412 (M + H)<sup>+</sup>.

#### Example 44

15 4-{[(2,6-Dichloro-phenyl)-hydroxy-methyl]-amino}-N-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide  
11 mg (0.02 mmol) of 2,6-dichloro-N-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-benzamide was dissolved in 2 ml tetrahydrofurane. Lithium aluminium  
hydride (6 mg, 0.16 mmol) was added to the reaction mixture under nitrogen  
20 atmosphere and solution was stirred overnight. The reaction mixture was filtered,  
evaporated to dryness and purified by on silica using gradient elution (chloroform  
to 4% methanol in chloroform) to obtain the title compound with a yield of 18%.  
MS (ESI<sup>+</sup>): m/z 475 (M + Na)<sup>+</sup>.

It will be appreciated that the methods of the present invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will be apparent for the specialist in the field that other embodiments exist and do not depart from the spirit of the invention. Thus, the described embodiments are  
5 illustrative and should not be construed as restrictive.

#### REFERENCES

- Cheng, Y., and Prusoff, W.H., 1973. *Biochem. Pharmacol.* 22: 3099
- Jasper, J.R., Lesnick, J.D., Chang, L.K., Yamanashi, S.S., Chang, T.C., Hsu, S.A.O., Daunt, D.A., Bonhaus, D.W., and Egen, R.M., 1998. *Biochem. Pharmacol.*  
10 55: 1035
- Regan, J.W., and Scheinin, M., 1992. *Biochem. Biophys. Acta* 1134: 169
- Marjamäki, A., Pihlavisto, M., Cockcroft, V., Heinonen, P., Savola, J.-M., and Scheinin, M., 1998. *Mol. Pharmacol.* 53: 370
- 15 Pohjanoksa, K., Jansson, C.C., Luomala, K., Marjamäki, A., Savola, J.-M., and Scheinin, M., 1997. *Eur. J. Pharmacol.* 35: 53
- Tian, W.-N., Duzic, E., Lanier, S.M., and Deth, R.C., 1993. *Mol. Pharmacol.* 45: 524
- Kumar, V.B., Reddy, M.V., *Indian J. Chem., Sect. B* (1985), 24B(12), 1298-1301
- 20 Farag, A.M., El-Mouafi, H.M., Khalifa, M., 1991. *Egypt. J. Pharm. Sci.* 32 (3-4), 951-9
- Wieland, T., and Jakobs, K.H., 1994. *Meth. Enzymol.* 237: 3


Heinonen et al.1999, The Journal of Clinical Endocrinology & Metabolism,  
84:2429

Link R E et al., 1996, Science 273:803

MacMillan L B et al., 1996, Science 273:801

## CLAIMS

1. A compound of formula (I)



or a pharmaceutically acceptable salt thereof wherein

- 5  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are independently of each other H, a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;
- $X$  is H, a straight or branched alkyl chain with 1 to 4 carbon atoms, phenyl or
- $Z$  is H, acetyl,  $-\text{CH}_2\text{-Ph-O-CF}_3$  or  $-\text{CH}_2\text{-Ph-CF}_3$ , where Ph is phenyl;
- 10  $Y$  is a ring structure optionally linked to formula (I) with an alkyl chain having one or two carbon atoms, wherein the ring structure is
- 15 a) phenyl optionally mono- or disubstituted and each substituent is independently selected from the group consisting of a halogen, a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, a halogen substituted methyl or methoxy group, a nitrile, an amide, amino, or a nitro group;
- 20 b) 2-benzimidazolyl, 2-imidazolyl, or 2- or 3-indolyl, wherein one N optionally has a substituent that is a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, or benzyl; and wherein the 2-benzimidazolyl, 2-imidazolyl, or 2- or 3-indolyl is optionally mono- or

disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

(c) pyridinyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen; or

(d) naphthyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

excluding 4-[(1*H*-benzimidazol-2-ylmethyl)-amino]-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-methyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide and *N*-(4-methyl-2-pyrimidinyl)-4-[(1*H*-benzimidazol-2-ylmethyl)-amino]-benzene-

2. A compound of formula (I) according to claim 1 wherein **R**<sub>1</sub> and **R**<sub>3</sub> are methyl and **R**<sub>2</sub>, **R**<sub>4</sub> and **R**<sub>5</sub> are H.

3. A compound of formula (I) according to claim 1 or 2 wherein **X** is H, **Y** is a phenyl optionally mono- or disubstituted with a straight or branched alkoxy group and **Z** is H.

4. A compound of formula (I) according to claim 2 and 3 wherein said phenyl is substituted and said alkoxy substituent is methoxy selected from the group consisting of 4-(2,4-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methoxybenzylamino)-benzenesulfonamide, 4-(3,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-


benzenesulfonamide, 4-(2,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide and *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methoxybenzylamino)-benzenesulfonamide.

5. A compound of formula (I) according to claim 1 or 2 wherein **X** is H ,  
5 **Y** is a phenyl optionally mono- or disubstituted with a straight or branched alkyl and/or a halogen and **Z** is H .

6. A compound of formula (I) according to claim 2 and 5 wherein said compound is selected from the group consisting of 4-benzylamino-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methylbenzylamino)-benzenesulfonamide,  
10 4-(2,4-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methylbenzylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, 4-(2,6-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide,  
15 4-(4-bromobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide and 4-(2,6-dichlorobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide.

7. A compound of formula (I) according to claim 2 selected from the group consisting of *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-indol-3-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-isobutyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(1-phenylethylamino)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[2-(2-methoxyphenyl)-ethylamino]-benzenesulfonamide and *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(naphthalen-2-ylmethyl)-amino]-benzenesulfonamide.  
25

## 8. Use of a compound of formula (I)



or a pharmaceutically acceptable salt thereof for the manufacture of a pharmaceutical preparation useful for the treatment or prevention of a disease mediated by the alpha-2B-adrenoceptor in a mammal wherein

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  are independently of each other H, a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

$X$  is H, a straight or branched alkyl chain with 1 to 4 carbon atoms, phenyl or -OH or =O.

10  $Z$  is H, acetyl,  $-\text{CH}_2\text{-Ph-O-CF}_3$  or  $-\text{CH}_2\text{-Ph-CF}_3$ , where Ph is phenyl;

$Y$  is a ring structure optionally linked to formula (I) with an alkyl chain having one or two carbon atoms, wherein the ring structure is

a) phenyl optionally mono- or disubstituted and each substituent is independently selected from the group consisting of a halogen, a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, a halogen substituted methyl or methoxy group, an acetyl, a nitrile, an amide, amino, or a nitro group;

b) 2-benzimidazolyl, 2-imidazolyl, or 2- or 3-indolyl, wherein one N optionally has a substituent that is a straight or branched alkyl or alkoxy chain with 1 to 4 carbon atoms, or benzyl; and wherein the 2-benzimidazolyl, 2-imidazolyl or 2- or 3-indolyl is optionally mono- or



disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

(c) pyridinyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen; or

(d) naphthyl optionally mono- or disubstituted and each substituent can independently be a straight or branched alkyl or alkoxy group with 1 to 4 carbon atoms, or a halogen;

excluding *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-methyl-1*H*-benzimidazol-2-ylmethyl)-amino] -benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-benzimidazol-2-ylmethyl)-amino] -benzenesulfonamide and *N*-[4-(4,6-dimethylpyrimidin-2-ylsulfamoyl)-phenyl]-4-ethoxy-benzamide.

according to claim 8 wherein  $R_1$  and  $R_3$  are methyl and  $R_2$ ,  $R_4$  and  $R_5$  are H.

10. Use of compound of formula (I) or a pharmaceutically acceptable salt thereof according to claim 8 or 9 wherein  $X$  is H,  $Y$  is a phenyl optionally mono- or disubstituted with a straight or branched alkoxy group and  $Z$  is H.

11. Use of compound of formula (I) or a pharmaceutically acceptable salt thereof according to claim 9 and 10 wherein said phenyl is substituted and said alkoxy substituent is methoxy and said compound is selected from the group consisting of 4-(2,4-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methoxybenzylamino)-benzenesulfonamide, 4-(3,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzene-

sulfonamide, 4-(2,5-dimethoxybenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide and *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methoxybenzylamino)-benzenesulfonamide.

12. Use of compound of formula (I) or a pharmaceutically acceptable salt thereof  
5 according to claim 8 or 9 wherein **X** is H, **Y** is a phenyl optionally mono- or disubstituted with a straight or branched alkyl and/or a halogen, and **Z** is H.

13. Use of compound of formula (I) or a pharmaceutically acceptable salt thereof  
according to claim 9 and 12 wherein said compound is selected from the group  
consisting of 4-benzylamino-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfonamide,  
10 *N*-(4,6-dimethylpyrimidin-2-yl)-4-(2-methylbenzylamino)-benzenesulfonamide,  
4-(2,4-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-yl)-benzenesulfon-  
amide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-(3-methylbenzylamino)-

benzenesulfonamide, 4-(2,5-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-2-  
15 yl)-benzenesulfonamide, 4-(2,6-dimethylbenzylamino)-*N*-(4,6-dimethylpyrimidin-  
2-yl)-benzenesulfonamide, 4-(4-bromobenzylamino)-*N*-(4,6-dimethylpyrimidin-2-  
yl)-benzenesulfonamide and 4-(2,6-dichlorobenzylamino)-*N*-(4,6-  
dimethylpyrimidin-2-yl)-benzenesulfonamide.

14. Use of a compound of formula (I) or a pharmaceutically acceptable salt  
20 thereof according to claim 9 wherein said compound is selected from the group  
consisting of 4-[(1*H*-benzimidazol-2-ylmethyl)-amino]-*N*-(4,6-dimethylpyrimidin-  
2-yl)-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-ethyl-1*H*-indol-  
3-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(1-  
isobutyl-1*H*-benzimidazol-2-ylmethyl)-amino]-benzenesulfonamide, *N*-(4,6-  
25 dimethylpyrimidin-2-yl)-4-(1-phenylethylamino)-benzenesulfonamide, *N*-(4,6-

dimethylpyrimidin-2-yl)-4-[2-(2-methoxyphenyl)-ethylamino]-benzenesulfonamide and *N*-(4,6-dimethylpyrimidin-2-yl)-4-[(naphthalen-2-ylmethyl)-amino]-benzenesulfonamide.

15. The use according to any of claims 8 to 14, wherein the disease is a coronary  
5 heart disease (CHD).

16. The use according to any of claims 8 to 14, wherein the disease is

- acute myocardial infarction (AMI),
- unstable angina pectoris,
- Prinzmetal's variant form of angina pectoris,
- 10 - other forms of chronic angina pectoris and CHD, or
- restenosis after coronary angioplasty.

17. The use according to any of claims 8 to 14, wherein the disease is essential hypertension.

18. The use according to any of claims 8 to 14, wherein the disease is a vascular  
15 disease, which is

vasoconstriction or hypoxic brain damage subsequent to subarachnoid  
haemorrhage,  
migraine,  
Raynaud's disease or cold intolerance,  
20 pre-eclampsia,  
male erectile dysfunction, or  
obesity.

19. The use according to any of claims 8 to 14, wherein said alpha-2B-adrenoceptor antagonist is administered to a mammal to potentiate the clinical efficacy of an anaesthetic and/or analgetic alpha-2-adrenoceptor agonist, said agonist not being selective for the alpha-2B-adrenoceptor subtype.
- 5 20. The use according to any of claims 8 to 14, wherein said alpha-2B-adrenoceptor antagonist is administered to an individual having a deletion of 3 glutamates from the glutamic acid repeat element of 12 glutamates (amino acids 297–309), in an acid stretch of 17 amino acids, located in the third intracellular loop of the receptor polypeptide.
- 10 21. The use according to claim 20 wherein said individual is a deletion/deletion genotype.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/FI 02/00643

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/69 C07D403/12 A61K31/635 A61P9/10 A61P9/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, CHEM ABS Data, EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 01 58454 A (OY JUVANTIA PHARMA LTD) 16 August 2001 (2001-08-16) the whole document	1-21
P,X	--- DATABASE STN INTERNATIONAL [Online] FILE CAPLUS; SANKYO COMPANY LTD: "Preparation of nitrophenylcarboxamide derivatives as non-viral gene delivery vectors, activated receptors	1-21
	retrieved from CAPLUS, accession no. 2001:816614 Database accession no. 135:357944 XP002902751 abstract & WO 01 83427 A (SANKYO COMPANY LTD) 8 November 2001 (2001-11-08) --- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

15 October 2002

Date of mailing of the international search report

01 11. 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gerd Strandell

## INTERNATIONAL SEARCH REPORT

onal Application No  
PCT/FI 02/00643

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE STN INTERNATIONAL [Online]  FILE CHEMCATS, order number  AE-848/12530187;  CAS Registry no. 380644-35-9,  1 July 2001 (2001-07-01)  "Benzenesulfonamide,  4-[1(1-methyl-1H-benzimidazol-2-yl)methyl]  amino]-N-(4-methyl-2-pyrimidinyl)-"  retrieved from CHEMCATS, accession no.  2002:1405436  XP002902752  abstract</p>	1
A	<p>---  US 5 292 740 A (BURRI KASPAR ET AL)  8 March 1994 (1994-03-08)  the whole document</p>	1-21
A	<p>---  RAJENDRA S. VARMA ET AL: "Potential  biologically active agents. XII. Synthesis  and antibacterial activity of substituted  p-methylamino-benzenesulfonamides."  JOURNAL INDIAN CHEM. SOC.,  vol. 55, no. 7, July 1978 (1978-07), pages  716-718, XP002902753  the whole document</p>	1-21
A	<p>---  DATABASE STN INTERNATIONAL [Online]</p>	1-21
A	<p>BEDENKO V. ET AL: "N4-acylsulfonamides. I.  Derivatives of  +-(2-ethoxybenzamido)benzenesulfonamide."  retrieved from CAPLUS, accession no.  1967:411468  Database accession no. 67:11468  XP002902754  abstract  &amp; CROAT. CHEM. ACTA.,  vol. 38, 1966, pages 309-311,  ---  US 6 150 389 A (MUNK S A ET AL)  21 November 2000 (2000-11-21)  the whole document  -----</p>	1-21

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No  
**PCT/FI 02/00643**

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 0158454	A	16-08-2001	AU	3551001 A		20-08-2001
			WO	0158454 A1		16-08-2001
			US	2002058618 A1		16-05-2002
-----						
WO 0183427	A	08-11-2001	AU	5261201 A		12-11-2001
			WO	0183427 A1		08-11-2001
-----						
US 5292740	A	08-03-1994	AT	197044 T		15-11-2000
			AU	653604 B2		06-10-1994
			AU	1812192 A		17-12-1992
			BG	60831 B2		30-04-1996
			CA	2071193 A1		14-12-1992
			CS	9201804 A3		16-12-1992
			DE	59209872 D1		23-11-2000
			DK	526708 T3		12-02-2001
			EE	3028 B1		15-08-1997
			EP	0526708 A1		10-02-1993
			ES	2152222 T3		01-02-2001
			FI	922746 A		14-12-1992
			HU	63152 A2		28-07-1993
			HU	9500261 A3		28-09-1995
			IE	921920 A1		16-12-1992
			IL	102138 A		12-09-1996
			JP	1997825 C		08-12-1995
			JP	5222003 A		31-08-1993
			JP	7030042 B		05-04-1995
			NO	922323 A		14-12-1992
			NZ	243074 A		25-11-2000
-----						
			RO	111268 B		30-08-1996
			SG	54209 A1		16-11-1998
			SK	279006 B6		06-05-1998
			BR	9202219 A		02-02-1993
			KR	235507 B1		15-12-1999
			MX	9202747 A1		31-05-1994
			RU	2086544 C1		10-08-1997
			ZA	9204126 A		24-02-1993
-----						
US 6150389	A	21-11-2000	US	5731337 A		24-03-1998
			US	6319935 B1		20-11-2001
			US	2002049241 A1		25-04-2002
			AU	687584 B2		26-02-1998
			AU	3007695 A		09-02-1996
			CA	2194702 A1		25-01-1996
			DE	69524223 D1		10-01-2002
			DE	69524223 T2		22-08-2002
			EP	0770068 A1		02-05-1997
			ES	2168373 T3		16-06-2002
			JP	10502643 T		10-03-1998
			WO	9601813 A1		25-01-1996
-----						



**This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record.**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**